

## REFERENCES

- Altun, Z., & Özdemir, M. (2018). The Role of the Music in Coping with Trauma Experiences. *European Journal of Education Studies*, 0. Retrieved from <https://oapub.org/edu/index.php/ejes/article/view/1853>
- Ames, N., Shuford, R., Yang, L., Moriyama, B., Frey, M., Wilson, F., Sundaramurthi, T., Gori, D., Mannes, A., Ranucci, A., Koziol, D., & Wallen, G.R. (2017). Music listening among postoperative patients in the intensive care unit: A randomized controlled trial with mixed-methods analysis. *Integrative Medicine Insights*, 12, 1-13. <https://doi.org/10.1177/1178633717716455>
- Anderson, C.A., Carnagey, N.L. & Eubanks, J. (2003). *Journal of Personality and Social Psychology*, 84(5), 960-971. <http://dx.doi.org/10.1037/0022-3514.84.5.960>
- Assagioli, R. (2000). *Psychosynthesis: a collection of basic writings*. Synthesis Center ed. Amherst, Mass.: Synthesis Center Inc. (in cooperation with the Berkshire Center for Psychosynthesis).
- Baker, F. & Bar, W. (2008). Can music preference indicate mental health status in young people? *Australasian Psychiatry*, 16(4), 284-288. <https://doi.org/10.1080/10398560701879589>
- Baird, A., & Samson, S. (2014). Music evoked autobiographical memory after severe acquired brain injury: Preliminary findings from a case series. *Neuropsychological Rehabilitation*, 24(1). <https://doi.org/10.1080/09602011.2013.858642>
- Behrman S., Chouliaras L., & Ebmeier K. P. (2014). Considering the senses in the diagnosis and management of dementia. *Maturitas*, 77(4), 305-310. <https://doi.org/10.1016/j.maturitas.2014.01.003>
- Bernardi, L., Porta, C., Cascucci, G., Balsamo, R., Bernardi, N. F., Fogari, R., & Sleight, P. (2009). Dynamic interactions between musical, cardiovascular, and cerebral rhythms in humans. *Circulation*, 119(225), 3171-3180. Retrieved from: <https://www.ncbi.nlm.nih.gov/pubmed/19569263>
- Bernardi, L., Porta, C., & Sleight, P. (2006). Cardiovascular, cerebrovascular, and respiratory changes induced by different types of music in musicians and non-musicians: the importance of silence. *Heart*, 92, 445–452. <http://dx.doi.org/10.1136/hrt.2005.064600>
- Bernatzky, G., Bernatzky, P., Hesse, H. P., Staffen, W., & Ladurner, G. (2004). Stimulating music increases motor coordination in patients afflicted with Morbus Parkinson, *Neuroscience Letters*, 361(1), 4-8. <http://dx.doi.org/10.1016/j.neulet.2003.12.022>

- Blais-Rochette, C. (2016). Music-evoked autobiographical memories, emotion regulation, time perspective, and mental health. *Musica Scientiae*, 20(1), 26–52. <https://doi.org/10.1177/1029864915626967>
- Carlson, E., Saarikallio, S., Toiviainen, P., Bogert, B., Kliuchko, M., & Brattico, E. (August 2015). Maladaptive and adaptive emotion regulation through music: A behavioral and neuroimaging study of males and females. *Frontiers in Human Neuroscience*. Retrieved from: <http://journal.frontiersin.org/article/10.3389/fnhum.2015.00466/full>
- Centers for Disease Control. (2018, January 23). Infection Control: Disinfection and Sterilization. Retrieved from: <https://www.cdc.gov/infectioncontrol/guidelines/disinfection/index.html>
- Chasin, M. (September 15, 2014). Programming hearing aids for listening and playing music, presented in partnership with the association of adult musicians with hearing loss (AAMHL). Retrieved from: <http://www.audiologyonline.com/articles/programming-hearing-aids-for-listening-12915>
- Chasin, M. (January 4, 2017). Back to basics: Music listening and hearing aids. are all approaches the same? stop and listen. Retrieved from: <http://www.hearingreview.com/2017/01/back-basics-music-listening-hearing-aids-approaches-stop-listen/>
- Clark M. E., Lipe A. W., & Bilbrey M. (1998). Use of music to decrease aggressive behaviors in people with dementia. *Journal of Gerontological Nursing*, 24, 10–17.
- Clements-Cortes, A., Pearson, C., & Chang, K. (2015). Creating effective music listening opportunities. Toronto, Ontario: Baycrest. Retrieved from <http://www.baycrest.org/care/culture-arts-innovation/therapeutic-arts/music-therapy/creating-effective-music-listening-opportunities>
- Croghan, N., Swanberg, A., Anderson M., & Arehart, K. (2015). Chosen listening levels for music with and without the use of hearing aids. *American Journal of Audiology*, 25, 161-166. [https://doi.org/10.1044/2016\\_AJA-15-0078](https://doi.org/10.1044/2016_AJA-15-0078)
- Dassa, A., & Amir, D. (2014). The role of singing familiar songs in encouraging conversation among people with middle to late stage Alzheimer's disease. *Journal of Music Therapy*, 51(2), 131-153. <https://doi.org/10.1093/jmt/thu007>
- DeDiego, A.C. (2019, January 25). The Use of Song Lyrics as an Expressive Arts Tool in Counseling: A Literature Review. *Ideas and Research You Can Use: VITAS 2015*, Article 84. Retrieved from: [https://www.counseling.org/docs/default-source/vistas/the-use-of-song-lyrics.pdf?sfvrsn=d3db432c\\_4](https://www.counseling.org/docs/default-source/vistas/the-use-of-song-lyrics.pdf?sfvrsn=d3db432c_4)
- Dingle, G., Kelly, P., Flynn, L., & Baker, F. (2015). The influence of music on emotions and cravings in clients in addiction treatment: A study of two clinical samples. *The Arts in Psychotherapy*, 45, 18-25. <https://doi.org/10.1016/j.aip.2015.05.005>

- Edworthy, J., & Waring, H. (2007). The effects of music tempo and loudness level on treadmill exercise. <https://doi.org/10.1080/00140130600899104>
- El Haj, M., Fassoti, L., & Allain, P. (2012). The involuntary nature of music-evoked autobiographical memories in Alzheimer's disease. *Consciousness and Cognition*, 21(1), 238-246. <https://doi.org/10.1016/j.concog.2011.12.005>
- Elcock, W. (February 9, 2016). How to change the speed of audio playback with these 4 android apps. *Guiding Technology*. Retrieved from: <https://www.guidingtech.com/56100/alter-speed-audio-playback-android-apps/>
- Epilepsy Society. (2012). *Musicogenic Epilepsy*. Retrieved from: <https://www.epilepsysociety.org.uk/music-and-epilepsy-.WaboQbpFw2w>
- Fligor, B. J., & Ives, T. (2006, October 19). *Does earphone type affect risk for recreational noise-induced hearing loss?* Retrieved from: <http://www.hearingconservation.org/docs/virtualpressRoom/FligorIves.pdf>
- First Steps Recovery. (2016, June 13). Why Music Can be A Trigger: Why Music Can Be a Drug Trigger in Sobriety. Retrieved from: <https://www.firststepsrecovery.com/why-music-can-be-a-trigger/>
- Fuld, G. L., Mulligan, D. A., Altmann, T. R., Brown, A., Christakis, D., Clarke-Pearson, K., ... Steiner, G. L. (2009). Policy statement - Impact of music, music lyrics, and music videos on children and youth. *Pediatrics*, 124(5), 1488-1494. <https://doi.org/10.1542/peds.2009-2145>
- Garrido, S., Eerola, T., & McFerran, K. (2017). Group Rumination: Social Interactions Around Music in People with Depression. *Frontiers in Psychology*, 8, 490. <https://doi.org/10.3389/fpsyg.2017.00490>
- Gerdner, L. A. (2012). Individualized music for dementia: Evolution and application of evidence-based protocol. *World Journal of Psychiatry*, 2(2), 26-32. <http://doi.org/10.5498/wjp.v2.i2.26>
- Gibbons, A. (1977). Popular music preferences of elderly people. *Journal of Music Therapy*, 14(4), 180-189. <https://doi.org/10.1093/jmt/14.4.180>
- Hamons, M., Jenkins, T., & Befi-Hensel, C. (2017). *Music, Memory, and Meaning*. Welk and Waters Publishing.
- Hanser, S. & Mandel, S. (2010). *Manage Your Stress and Pain through Music*. Berklee Press.
- HermanMiller Healthcare Research Summary (2006). Sound practices: Noise control in the healthcare environment. Retrieved from:

[https://www.hermanmiller.com/MarketFacingTech/hmc/research\\_summaries/pdfs/wp\\_Sound\\_Practices.pdf](https://www.hermanmiller.com/MarketFacingTech/hmc/research_summaries/pdfs/wp_Sound_Practices.pdf)

- Hicks, D. (2016). *Separating myth from reality: Cleaning and disinfecting mobile devices*. Retrieved from: [https://www.zebra.com/content/dam/zebra\\_new\\_ia/en-us/solutions-verticals/vertical-solutions/healthcare/white-paper/cleaning-disinfecting-mobile-devices-hc-en-us-global.pdf](https://www.zebra.com/content/dam/zebra_new_ia/en-us/solutions-verticals/vertical-solutions/healthcare/white-paper/cleaning-disinfecting-mobile-devices-hc-en-us-global.pdf)
- Janata, P. (2009). The Neural Architecture of Music Evoked Autobiographical Memories. *Cerebral Cortex*, 12, 2579-2594. <https://doi.org/10.1093/cercor/bhp008>
- Janelli L. M., Kanski G. W., & Wu Y. W. (2002). Individualized music--a different approach to the restraint issue. *Rehabilitation Nursing*, 27, 221–226.
- Jeppesen, E., Pedersen, C.M., Larsen, K.R, Walsted, E.S., Rehl, A., Ehrenreich, J., Schnoor, S., & Backer, V. (2019). Listening to music prior to bronchoscopy reduces anxiety--a randomised controlled trial. *European Clinical Respiratory Journal*, 6(1). doi: <https://doi.org/10.1080/20018525.2019.1583517>Jespersen, K. V., Koenig, J., Jennum, P., & Vuust, P. (2015). Music for insomnia in adults. Retrieved from: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD010459.pub2/full>
- Johnson, L., Deatrick, E. J., & Oriol, K. (2012). The use of music to improve exercise participation in people with dementia: A pilot study. *Physical & Occupational Therapy*, 30(2). <https://doi.org/10.3109/02703181.2012.680008>
- Josefczyk, J. (September 19, 2017). Listening programs: Fast track to nowhere fast. *Music Therapy and End-of-Life Care Blog*. Retrieved from: <https://musictherapyeolcare.com/2017/09/19/listening-programs-fast-track-to-nowhere-fast/>
- Kampfe, J., Sedlmeirer, P., & Renkewitz, F. (2011). The impact of background music on adult listeners: A meta-analysis. *Psychology of Music*, 39(4), 424-448. <https://doi.org/10.1177/0305735610376261>
- Karageorghis, C. I., & Priest, D. L. (July 7, 2008). Music in sport and exercise : An update on research and application. *The Sport Journal*, 19. Retrieved from: <http://thesportjournal.org/article/music-sport-and-exercise-update-research-and-application/>
- Kardous, C., Themann, C. L., Morata, T. C., & Lotz, W. G. (February 8, 2016). Understanding noise exposure limits: Occupational vs. general environmental noise. Retrieved from <https://blogs.cdc.gov/niosh-science-blog/2016/02/08/noise/>
- Keith, D.R., Russel, K., & Weaver, B.S. (2009). The Effects of Music Listening on Inconsolable Crying in Premature Infants. *Journal of Music Therapy*, 46(3), 191-203. <https://doi.org/10.1093/jmt/46.3.191>

- Koelsch, S. (2010). Towards a neural basis of music-evoked emotions. *Trends in Cognitive Sciences*, 14(3), 131–137. <https://doi.org/10.1016/j.tics.2010.01.002>
- Labbé, E., Schmidt, N., Babin, J., & Pharr, M. (2007). Coping with stress: The effectiveness of different types of music. *Applied Psychophysiological Biofeedback*, 32, 163-168. <https://doi.org/10.1007/s10484-007-9043-9>
- Lamparero-Plokhotnikov, J. L. (2015). *The use of music in dementia care: A literature study*. Degree Thesis in Health Care and Social Welfare, Vasa Education: Nurse, Bachelor of Health Care Vasa. (Unpublished master's thesis). Vasa Education.
- Leblanc, A. (1982). An interactive theory of music preference. *Journal of Music Therapy*, 19(1), 28-45. <https://doi.org/10.1093/jmt/19.1.28>
- Lesiuk, T. (2010). The effect of preferred music on mood and performance in a high cognitive demand occupation. *Journal of Music Therapy*, 47(2), 137-154. <https://doi.org/10.1093/jmt/47.2.137>
- Levitin, D. (2009, May 21). Music helps scientists understand brain function [Interview]. Retrieved from: <https://www.pbs.org/wnet/musicinstinct/mi-blog/interview/interview-with-daniel-levitin-part-1/18/>
- Loomba, R. S., Arora, R., Shah, P. H., Chandrasekar, S., & Molnar, J. (2012). Effects of music on systolic blood pressure, diastolic blood pressure, and heart rate: a meta-analysis. *Indian Heart Journal*, 64(3), 309–313. [https://doi.org/10.1016/S0019-4832\(12\)60094-7](https://doi.org/10.1016/S0019-4832(12)60094-7)
- Macnay, S. (1995). The influence of preferred music on the perceived exertion, mood, and time estimation scores of patients participating in a cardiac rehabilitation exercise program. *Music Therapy Perspectives*, 13, 91-96. <https://doi.org/10.1093/mtp/13.2.91>
- Marks, R., & Landaira, M. (2015). Musical exercise: A novel strategy for advancing healthy aging. *Healthy Aging Research*, 4(31), 1-8. doi:10.12715/har.2015.4.31
- Martin-Saavedra, JS., Vergara-Mendez, LD., Pradilla, I., Velez-van-Meerbeke, A., Talero-Gutierrez, C. (2018). Standardizing music characteristics for the management of pain: A systematic review and meta-analysis of clinical trials. *Complementary Therapies in Medicine*, 41, 81-89. <https://doi.org/10.1016/j.ctim.2018.07.008>
- Mitchell, L., & MacDonald, R. (2006). An experimental investigation of the effects of preferred and relaxing music listening on pain perception. *Journal of Music Therapy*, 43(4), 295-316. <https://doi.org/10.1093/jmt/43.4.295>
- Mohney, B. (2015). *The effect of client preferred music on workout efficiency and perceived energy exertion among collegiate football players* (Unpublished master's thesis). Florida State.

- Nilsson, U. (2009). Soothing music can increase oxytocin levels during bed rest after open-heart surgery: A randomised control trial. *Journal of Clinical Nursing*, 18(15).  
<https://doi.org/10.1111/j.1365-2702.2008.02718.x>
- Papinczak, Z.E., Dingle, G.A., Stoyanov, S.R., Hides, L., & Zelenko, O. (2015). Young people's uses of music for well-being. *Journal of Youth Studies*, 18(9), 1119-1134.  
<https://doi.org/10.1080/13676261.2015.1020935>
- Pedersen, S., Andersen, P., Lugo, R., Andreassen, M., & Sütterlin, S. (2017). Effects of music on agitation in dementia: A meta-analysis. *Frontiers in Psychology*, 8(742).  
<https://doi.org/10.3389/fpsyg.2017.00742>
- Portnuff, C., & Fligor, B. (October 17, 2006). Researchers recommend safe listening levels for Apple iPod. Retrieved from: <https://phys.org/news/2006-10-safe-apple-ipod.html>
- Proudfoot, J., Whitton, A., Parker, G., Doran, J., Manicavasagar, V., & Delmas, K. (2012). Triggers of mania and depression in young adults with bipolar disorder. *Journal of Affective Disorders*, 143(1-3), 196-202. <https://doi.org/10.1016/j.jad.2012.05.052>
- Ransom, P.F. (2015). Message in the Music: Do Lyrics Influence Well-Being? Master of Applied Positive Psychology (MAPP) Capstone Projects. 94.  
[http://repository.upenn.edu/mapp\\_capstone/94](http://repository.upenn.edu/mapp_capstone/94)
- Särkämö, T., Ripollés, P., Vepsäläinen, H., Autti, T., Silvennoinen, H. M., Salli, E., & Rodríguez-Fornells, A. (2014). Structural changes induced by daily music listening in the recovering brain after Middle Cerebral Artery Stroke: A voxel-based morphometry study. *Frontiers in Human Neuroscience*, 8, 245. <https://doi.org/10.3389/fnhum.2014.00245>
- Särkämö T., Tervaniemi M., Laitinen S., Forsblom A., Soynila S., Mikkonen M., & Hietanen M. (2008). Music listening enhances cognitive recovery and mood after middle cerebral artery stroke, *Brain*, 131(3), 866–876. <https://doi.org/10.1093/brain/awn013>
- Schaefer, R.S. (2014). Auditory rhythmic cueing in movement rehabilitation: Findings and possible mechanisms. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 369: 20130402. <https://doi.org/10.1098/rstb.2013.0402>
- Sena Moore, K. (2014, August 15). 5 Problems Music Can Create [Blog post]. Retrieved from: <https://www.psychologytoday.com/us/blog/your-musical-self/201408/5-problems-music-can-create>
- Sena Moore, K. (2016). Understanding the Influence of Music on Emotions: A Historical Review. *Music Therapy Perspectives*, 35(2), 131-143. <https://doi.org/10.1093/mtp/miw026>
- Shoemark H., & Dearn, T. (2016). Music therapy in the medical care of infants. *The Oxford Handbook of Music Therapy*. (Ed.) Jane Edwards: New York, NY, p. 24.

- Standley, J.M. (2002). A meta-analysis of the efficacy of music therapy for premature infants. *Journal of Pediatric Nursing*, 17(2), 107-113. <https://doi.org/10.1053/jpdn.2002.124128>
- Sung, H. C., Chang, A. M., & Lee, W. L. (2010). A preferred music listening intervention to reduce anxiety in older adults with dementia in nursing homes. *Journal of Clinical Nursing*, 19(7-8), 1056-1064. <https://doi.org/10.1111/j.1365-2702.2009.03016.x>
- Takahashi, T., & Matsushita, H. (2006). Long-term effects of music therapy on elderly with moderate/severe dementia. *Journal of Music Therapy*, 43, 317-333. <https://doi.org/10.1093/jmt/43.4.317>
- Tan, Y. Z., Ozdemir, S., Temiz, A., & Celik, F. (2015). The effect of relaxing music on heart rate and heart rate variability during ECG GATED-myocardial perfusion scintigraphy. *Complementary Therapies in Clinical Practice*, 21(2), 137-140. <https://doi.org/10.1016/j.ctcp.2014.12.003>
- Taylor, D. B. (2010). *Biomedical Foundations of Music as Therapy*, 2<sup>nd</sup> ed. Eau Claire: Barton Publications.
- Thomas D. W., Heitman R. J., & Alexander T. (1997). The effects of music on bathing cooperation for residents with dementia. *Journal of Music Therapy*, 34, 246-259. <https://doi.org/10.1093/jmt/34.4.246>
- Thomas, D. W., & Smith, M. (2009). The effect of music on caloric consumption among nursing home residents with dementia of the Alzheimer's type. *Activities, Adaptation & Aging*, 33(1). <https://doi.org/10.1080/01924780902718566>
- To, L. (2016). Exploring the effect of sound and music on health in hospital settings: A narrative review. *International Journal of Nursing Studies*, 63, 82-100. <https://doi.org/10.1016/j.ijnurstu.2016.08.008>
- University of California-Davis. (2009, February 24). Brain Hub That Links Music, Memory and Emotion Discovered. *ScienceDaily*. Retrieved from: <http://www.sciencedaily.com/releases/2009/02/090223221230.htm>
- Vachiramam, V., Sobanko, J., Rattanaumpawan, P., & Miller, C.J. (2013). Music reduces patient anxiety during MOHS surgery: An open-label randomized controlled trial. *Dermatological Surgery*, 39(2), 298-305. <https://doi.org/10.1111/dsu.12047>
- Vaisberg, J.M., Folkeard, P., Parsa, V., Froehlich, M., Littmann, V., Macpherson, E.A., & Scollie, S. (2017, August). Comparison of music sound quality between hearing aids and music programs. *AudiologyOnline*, Article 20872. Retrieved from <http://www.audiologyonline.com>
- Walworth, D. (2003). The effect of preferred music genre selection versus preferred song selection on experimentally induced anxiety levels. *Journal of Music Therapy*, 40(1), 2-14. <https://doi.org/10.1093/jmt/40.1.2>

Warren, M. (December 19, 2016). The impact of music therapy on mental health. *National Alliance on Mental Health*. Retrieved from: <https://www.nami.org/Blogs/NAMI-Blog/December-2016/The-Impact-of-Music-Therapy-on-Mental-Health>

Weisman-Fogel, I., Granovsky, Y., & Bar-Shalita, T. (2017). Sensory over-responsiveness among healthy subjects is associated with a pro-nociceptive state. *Pain Practice*, 18(4), 473-486. <https://doi.org/10.1111/papr.12619>

World Health Organization. (2017). Make listening safe. Retrieved from: <http://www.who.int/pbd/deafness/activities/MLS/en/>